

Introduction

Do heavy quarks contribute to the proton PDFs at low scales?

We look at the charm PDF in NNPDF4.0 [1]:


- The charm PDF is parametrized as an independent combination at scale $Q = 1.65 \text{ GeV}$, and $n_f = 4$ (4FNS):

$$xc^+(x, Q_0, \Theta) = \frac{1}{4} \left(x^{\alpha_\Sigma} (1-x)^{\beta_\Sigma} NN_{\Sigma}(x, \Theta) - x^{\alpha_{T15}} (1-x)^{\beta_{T15}} NN_{T15}(x, \Theta) \right) \quad (1)$$

- We always assume: $c(x, Q) = \bar{c}(x, Q) \rightarrow c^- = 0$.
- At the fitting scale, we observe that constraints are coming mainly from collider data. NNPDF4.0 is consistent with EMC data.


Fitted or Perturbative?

- Perturbative charm (PC)** functional form is fully determined by the DGLAP evolution and the initial boundary conditions on gluons and light quarks.




- PDF uncertainties are clearly not the dominant source of uncertainties. Needs to **estimate MHOU**. PC is also highly correlated to the mass value.
- Fitted charm is favored by data: $\chi^2_{fitted\ ch} = 1.17$, $\chi^2_{pert\ ch} = 1.19$ mainly due to a worsening of the LHC W, Z and top pair data sets.
- Fully perturbative charm is not compatible with the fitted one especially at large x, even if MHOU (N³LO - NNLO) are considered.

From 4FNS to 3FNS




The Intrinsic Charm

- Below the charm mass scale the perturbative charm is vanishing by definition.
- Fitted charm in 4FNS contains both the intrinsic and the perturbative components.
- To obtain the **Intrinsic charm (IC)** we start from the fitting scale we evolve the NNPDF4.0 baseline to $Q = m_c = 1.51 \text{ GeV}$. When passing the heavy quark threshold we need to invert the matching conditions.
- The remaining part of the charm PDF is the intrinsic component, which is **scale independent**



IC stability


- IC valence-like peak at large-x is a stable feature.
- IC vanishes for $x \rightarrow 0$. In this region the effect of MHOU is large.
- IC is stable upon mass variation in the range $m_c = 1.51 \pm 0.13 \text{ GeV}$.
- The momentum fraction carried by the IC component is: $0.62 \pm (0.28_{pdf} \pm 0.54_{mhou})\%$
- In our best estimation (NNPDF4.0+EMC+LHCb Zc) we reach a 3σ local evidence.



Comparison with models

The determination of the IC can be compared to some non perturbative models:


- BHPS** [2]: the proton fluctuates to a pentaquark state $p \rightarrow u\bar{u}dc\bar{c}$. It assumes $c = \bar{c}$.
- Meson Baryon** model [3]: fluctuations to a charmed baryon plus meson state (ex: $p \rightarrow \lambda_c^+ + \bar{D}_0$). It can account for c, \bar{c} asymmetry.



Reasonable agreement is found with both models. However, the analysis is not yet conclusive in this respect and there are still some limitations (ex: is $c^- \neq 0$?, mass corrections)

Impact on LHC observables

We validate our observation of IC testing the impact on prediction for: **Z + c production at LHCb** [4]



- Better agreement is found with the NNPDF4.0 fitted charm especially in the **forward region**
- High correlation with the charm PDF and LHCb observable.
- Predictions are also stable upon charm mass variation

References

- R. D. Ball et al., *The Path to Proton Structure at One-Percent Accuracy*, [arXiv:2109.02653](https://arxiv.org/abs/2109.02653).
- S. Brodsky, P. Hoyer, C. Peterson, and N. Sakai, *The intrinsic charm of the proton*, *Physics Letters B* **93** (1980), no. 4, 451–455.
- T. J. Hobbs, J. T. Londergan, and W. Melnitchouk, *Phenomenology of nonperturbative charm in the nucleon*, *Phys. Rev. D* **89** (2014), no. 7, 074008, [[arXiv:1311.1578](https://arxiv.org/abs/1311.1578)].
- LHCb** Collaboration, R. Aaij et al., *Study of Z Bosons Produced in Association with Charm in the Forward Region*, *Phys. Rev. Lett.* **128** (2022), no. 8, 082001, [[arXiv:2109.08084](https://arxiv.org/abs/2109.08084)].